



Technical Data

Toluidine Blue (0.1gm per vial)

FD051

This supplement is recommended for detection of deoxyribonuclease activity.

Composition

Per vial sufficient for 1000 ml medium

Ingredients

Toluidine blue

Concentration

0.1g

Directions:

Rehydrate the contents of 1 vial in 5 ml of distilled water and add in 1000 ml of rehydrated DNase Test Agar Base [M482](#) / DNase Test HiVeg™ Agar Base [MV482](#) . Heat to boiling to dissolve the medium completely. Sterilize by autoclaving the medium at 15 lbs pressure (121°C) for 15 minutes. Cool and pour in sterile petri plates. Alternatively 0.1% aqueous solution of toluidine blue can be used to flood the growth on DNase Test Agar plates.

Storage and Shelf Life

Store between 10-30°C. Use before expiry date on the label.

Revision : 1 / 2012

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.