

Technical Data

Tryptone Glucose Extract Agar (Tryptone Glucose Yeast Extract Agar)

M014

Tryptone Glucose Yeast Extract Agar is recommended for enumeration of bacteria in water, air, milk and dairy products.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	5.000
Yeast extract	3.000
Glucose	1.000
Agar	15.000
Final pH (at 25°C)	7.0±0.2

^{**}Formula adjusted, standardized to suit performance parameters

Directions

Suspend 24 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Tryptone Glucose Yeast Extract Agar was originally developed by Bowers and Hucker (1) which they called as Tryptone Glucose Skim Milk Agar. Later on it was modified to the present composition for the cultivation and enumeration of bacteria in air, water (2), milk and dairy products (3). Various authors have studied different aspects of this medium like study of thermophilic bacteria in milk (4), influence of incubation temperature (5) etc. It is used as a standard medium for the bacteriological plate count of milk and dairy products (6).

Casein enzymic hydrolysate, yeast extract provide nitrogenous compounds, vitamin B complex and other essential growth nutrients. Glucose is the energy source. For the enumeration purposes, pour plate method is suggested. Medium must be quickly poured into Petri dishes if milk sample is to be tested, because the milk may get flocculated if the medium remains hot for longer period of time.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light yellow coloured Clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 2.4% w/v aqueous solution at 25°C. pH: 7.0±0.2

pН

6.80-7.20

Cultural Response

M014: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery
Cultural Response			
Bacillus subtilis ATCC 6633	50-100	good-luxuriant	>=70%
Enterobacter aerogenes	50-100	good-luxuriant	>=70%
ATCC 13048			
Escherichia coli ATCC	50-100	good-luxuriant	>=70%
25922		-	

HiMedia Laboratories Technical Data

Enterococcus faecalis ATCC 29212	C 50-100	good-luxuriant >=70%
Lactobacillus casei ATCC 9595	50-100	good-luxuriant >=70%
Pseudomonas aeruginosa ATCC 27853	50-100	good-luxuriant >=70%
Staphylococcus aureus	50-100	good-luxuriant >=70%

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label..

Reference

- 1. Bowers and Hucker, 1935, Tech. Bull., 228, N.Y.State Agr. Expt. Station.
- 2. Eaton A. D., Clesceri L. S. and Greenberg A. W., (Eds.), 2005, Standard Methods for the Examination of Water and Wastewater, 21st Ed., APHA, Washington, D.C.
- 3. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.
- 4. Pickett, 1928, Tech. Bull. 147, N.Y. State Agr. Expt. Station.
- 5. Dennis and Weiser, 1937, J.Dairy Science, 20: 445.
- 6. Abele C. A., Am. J. Pub. Health, 1939, 29: 821.

Revision: 02 / 2015

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMediaTM publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMediaTM Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.