



Czapek Dox Agar

M075

Intended use

Czapek Dox Agar is a semisynthetic medium used for the general cultivation of fungi from water samples.

Composition**

Ingredients	Gms / Litre
Sucrose	30.000
Sodium nitrate	2.000
Dipotassium phosphate	1.000
Magnesium sulphate	0.500
Potassium chloride	0.500
Ferrous sulphate	0.010
Agar	15.000
Final pH (at 25°C)	7.3±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 49.01 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

Fungi, including yeasts and filamentous species or moulds are ubiquitously distributed in nature. Czapek Dox Agar is a semi-synthetic medium used for the cultivation of fungi, containing sodium nitrate as the sole source of nitrogen. This medium is prepared according to the formula developed by Thom and Church (4), which has a defined chemical composition. Czapek Dox Agar is recommended by APHA (1) for isolation of *Aspergillus*, *Penicillium*, *Paecilomyces* and some other fungi with similar physiological requirements.

Sucrose serves as the sole source of carbon while sodium nitrate serves as the sole source of nitrogen. Dipotassium phosphate buffers the medium. Magnesium sulphate, potassium chloride, ferrous sulphate serves as sources of essential ions.

Type of specimen

Water samples.

Specimen Collection and Handling:

For water samples, follow appropriate techniques for sample collection, processing as per guidelines and local standards.(1) After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1.This medium is general purpose medium and may not support the growth of fastidious organisms.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Please refer disclaimer Overleaf.

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light yellow coloured, clear to slightly opalescent gel with a slight precipitate forms in Petri plates

Reaction

Reaction of 4.9% w/v aqueous solution at 25°C. pH : 7.3±0.2

pH

7.10-7.50

Cultural Response

M075: Cultural characteristics observed after an incubation at 25-30°C for 48-72 hours.

Organism	Growth	Inoculum (CFU)	Recovery
<i>Aspergillus brasiliensis</i> ATCC 16404 (00053*)	luxuriant		>=50%
<i>Candida albicans</i> ATCC 10231 (00054*)	luxuriant	50-100	>=50%
<i>Saccharomyces cerevisiae</i> ATCC 9763	luxuriant	50-100	>=50%

Key : * -Corresponding WDCM numbers

Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 20-30°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (2,3).

References

1. Baird R.B., Eaton A.D., and Rice E.W., (Eds.), 2015, Standard Methods for the Examination of Water and Wastewater, 23rd ed., APHA, Washington, D.C.
2. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
3. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock, D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
4. Thom and Church, 1926, The Aspergilli, 39.

Revision : 03 / 2018

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.