



## Nutrient Agar, pH 6.0 with 0.8% NaCl

M090

Nutrient Agar, pH 6.0 with 0.8% NaCl is used for cultivation of bacteria requiring slightly acidic pH.

### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Beef extract	3.000
Sodium chloride	8.000
Agar	15.000
Final pH ( at 25°C)	6.0±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 31 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

### Principle And Interpretation

Nutrient Media are general purpose media used for the examination of water and dairy products according to Standard Methods for the Examination of Water and Waste water (1) and Dairy Products (2). Nutrient Agar, pH 6.0 with 0.8% NaCl is a modification of Nutrient Agar w/ 0.8 % NaCl and recommended by APHA (3). In the former, the pH of the medium is adjusted to 6.0 to allow the growth of organisms requiring slightly acidic pH. Since the medium contains 0.8 % sodium chloride, it can be used as a base for enrichment with blood or ascetic fluid or other supplements for cultivation of fastidious microorganisms. Sodium chloride maintains the osmotic balance so that red blood cells do not rupture when blood is added as supplement (1).

Beef extract and peptic digest of animal tissue provide the necessary nitrogen compounds, carbon, vitamins and also some trace ingredients to the nonfastidious organisms like *Bacillus subtilis* and *Staphylococcus aureus* . Sodium chloride maintains osmotic equilibrium of the medium.

### Quality Control

#### Appearance

Cream to yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 1.5% Agar gel

#### Colour and Clarity of prepared medium

Light yellow to amber coloured clear to slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 3.1% w/v aqueous solution at 25°C. pH : 6.0±0.2

#### pH

5.80-6.20

#### Cultural Response

M090: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery
----------	----------------	--------	----------

#### Cultural Response

<i>Bacillus subtilis</i> ATCC 6633	50-100	good	50-70%
<i>Candida albicans</i> ATCC 10231	50-100	luxuriant	>=70%
<i>Staphylococcus aureus</i> ATCC 25923	50-100	good	50-70%

## Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

## Reference

1. Clesceri L. S, Greenberg A. E. and Eaton A. D., (Eds.), 1998, Standard Methods for the Examination of Water and Wastewater, 20th Ed., APHA, Washington, D.C.
2. American Public Health Association, 1978, Standard Methods for the Examination of Dairy Products, 14th Ed., APHA, Inc., Washington, D.C.
3. Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., American Public Health Association, Washington, D.C.

Revision : 2 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.