



## MIU Medium Base

M1076

MIU Medium Base is recommended for detection of motility, urease and indole production.

### Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	10.000
Dextrose	1.000
Sodium chloride	5.000
Phenol red	0.010
Agar	2.000
Final pH ( at 25°C)	6.8±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 18 grams in 950 ml distilled water. Heat to boiling to dissolve the medium completely. Dispense in 95 ml amounts into flasks and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to about 50-55°C and add aseptically 5 ml sterile 40% Urea solution (FD048) per 95 ml basal medium. Mix well and dispense into sterile test tubes. Allow to cool in an upright position.

### Principle And Interpretation

MIU Medium Base is formulated to detect motility, urease and indole production in single tube.

Casein enzymic hydrolysate provide amino acids and other nitrogenous substances. Sodium chloride maintains osmotic equilibrium. Dextrose is fermentable carbohydrate. Phenol red is the pH indicator which turns pink- red in alkaline conditions. The test cultures are stab-inoculated.

Motility and urease reactions are read before testing Indole production. Motile organisms show either diffused growth or turbidity extending away from stab inoculation line while nonmotile organisms grow along the stabline. Organisms that utilize urea, produce ammonia which makes the medium alkaline, showing pink-red colour by change in the phenol red indicator (1). Indole is produced from tryptophan present in casein enzymic hydrolysate (2,3). The indole produced combines with the aldehyde present in the Kovac's reagent to form a red complex.

### Quality Control

#### Appearance

Light orange to light pink coloured homogeneous free flowing powder

#### Gelling

Semisolid, comparable with 0.2% Agar gel.

#### Colour and Clarity of prepared medium

Yellowish orange coloured clear to slightly opalescent gel is obtained in tubes as butts after addition of urea solution.

#### Reaction

Reaction of basal medium (1.8 gm suspended in 95 ml distilled water) at 25°C. pH : 6.8±0.2

#### pH

6.60-7.00

#### Cultural Response

M1076: Cultural characteristics observed with added 40% Urea solution (FD048) after an incubation at 35-37°C for 18 - 24 hours.

Organism	Growth	Indole	Motility	Urease activity
<b>Cultural Response</b> <i>Escherichia coli</i> ATCC 25922	luxuriant	Positive reaction, red ring at the	Positive, growth away from stabline	Negative reaction, no change

<i>Klebsiella pneumoniae</i> ATCC 13883	luxuriant	interface of the medium Negative reaction no colour development / cloudy ring	causing turbidity Negative growth along the stabline, surrounding medium remains clear	Weakly positive
<i>Proteus mirabilis</i> ATCC 25933	luxuriant	Negative reaction no colour development / cloudy ring	Positive, growth away from stabline causing turbidity	Positive reaction, cerise colour
<i>Proteus vulgaris</i> ATCC 13315	luxuriant	Positive reaction, red ring at the interface of the medium	Positive, growth away from stabline causing turbidity	Positive reaction, cerise colour
<i>Salmonella typhimurium</i> ATCC 14028	luxuriant	Negative reaction no colour development / cloudy ring	Positive, growth away from stabline causing turbidity	Negative reaction, no change

### Storage and Shelf Life

Store below 30°C in tightly closed container and use freshly prepared medium . Use before expiry date on the label.

### Reference

1. Rustigian and Stuart (1941) Proc. Soc. Exp. Biol. Med., 47:108.
2. McFaddin J.F. (1985) Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore
3. Ewing (1986) Edwards and Ewings 'Identification of Enterobacteriaceae', 4th ed. Elsevier Science Publishing Co., Inc., New York.

Revision : 2 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.