



## M-FC Agar Base, Modified

M1124

M-FC Agar Base, Modified is used for rapid enumeration of *Klebsiella* using membrane filter technique.

### Composition\*\*

Ingredients	Gms / Litre
Tryptose	10.000
Proteose peptone	5.000
Yeast extract	3.000
Sodium chloride	5.000
Inositol	10.000
Bile salts mixture	1.500
Aniline blue	0.100
Agar	15.000
Final pH ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 49.6 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. DO NOT AUTOCLAVE. Add 10 ml of 1% Rosolic Acid (FD058). Cool below 45°C and add 50 mg Carbenicillin. Mix well and pour into sterile Petri plates.

### Principle And Interpretation

M-FC Agar Base, Modified is used for the enumeration of *Klebsiella* using membrane filter technique. *Klebsiella* are widely distributed in nature, occurring in soil, water, grains, vegetation etc. Wood pulp, paper mills, textile finishing plants and sugarcane processing operations contain large numbers of *Klebsiella* in their effluents and are often in the predominant coliform in such effluents. M-FC Agar, Modified is formulated as per APHA (1) for enumeration of *Klebsiella*. M-FC Agar is modified by replacing lactose by inositol and adding Carbenicillin.

Proteose peptone, tryptose and yeast extract in the medium provide necessary nutrients for the growth of faecal coliforms. Inositol is the fermentable carbohydrate and the carbon source in the medium. Bile salts mixture inhibits the growth of contaminating gram-positive microorganisms. Aniline blue is a triphenyl methane dye, which suppresses the growth of many gram-positive microorganisms. Also, along with rosolic acid it forms the indicator system in the medium. Carbenicillin inhibits accompanying coliforms and other bacteria and helps in selective isolation of *Klebsiella* species.

Sample volume is selected to yield 20 to 60 *Klebsiella* colonies per membrane. This membrane filter is aseptically placed on agar surface. Occasional false positive results may be occurred due to *Enterobacter* species. *Klebsiella* colonies appear deep blue to blue grey due to aniline blue present in the medium. *Klebsiella* colonies will form blue or bluish grey coloured. Presumptive colonies should be further confirmed by performing the biochemical tests.

### Quality Control

#### Appearance

Light yellow to greyish yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 1.5% Agar gel

#### Colour and Clarity of prepared medium

After Addition of 1% Rosolic Acid : Red coloured clear to slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 4.96% w/v aqueous solution at 25°C. pH : 7.4±0.2

#### pH

7.20-7.60

### Cultural Response

M1124: Cultural characteristics observed with added 1% Rosolic Acid (FD058) after an incubation at 35 - 37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Colour of colony (on membrane filter)
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	good-luxuriant	pink or occasionally pale yellow
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	good-luxuriant	deep blue to blue grey

### Storage and Shelf Life

Store below 30°C in tightly closed container and use freshly prepared medium. Use before expiry date on the label.

### Reference

1. Eaton A. D., Clesceri L. S. and Greenberg A. W., (Eds.), 2005, Standard Methods for the Examination of Water and Wastewater, 21st Ed., APHA, Washington, D.C.

Revision : 2 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.