



## Bromo Cresol Purple Broth w/Dextrose

M1463

Bromo Cresol Purple Broth with Dextrose is used for identification of *Escherichia coli* and coliform bacteria from water samples.

### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	10.000
Meat extract	3.000
Sodium chloride	5.000
Dextrose	10.000
Bromo cresol purple	0.020
Final pH ( at 25°C)	7.2±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 28.02 grams in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Dispense in tubes containing inverted Durham's tubes and sterilize by autoclaving at 115°C for 20 minutes.

### Principle And Interpretation

The coliform group of bacteria is the principal indicator of suitability of water for domestic, industrial or other uses. The coliform group density has been established as a criterion of the degree of pollution and thus of sanitary quality. Faecal Streptococci and Enterococci are also indicators of faecal pollution (1). Where it is claimed that drinking water has been processed for safety, the finding of such organism demonstrates a failure of the process. It is a valuable bacterial indicator for determining the extent of fecal contamination of recreational surface waters or drinking water (2).

Bromo Cresol Purple Broth with Dextrose is used for the identification of *Escherichia coli* and coliforms from water. It is used for enrichment and determining the titre of coliforms in the bacteriological analysis of drinking water (3, 4).

The medium contains peptic digest of animal tissue and meat extract, which supplies the essential nutrients for *E. coli* and other coliforms. Sodium chloride maintains the osmotic equilibrium of the medium. Dextrose upon fermentation by coliforms produce acid and is indicated by the pH indicator bromo cresol purple. It turns yellow at acidic pH.

### Quality Control

#### Appearance

Cream to yellow homogeneous free flowing powder

#### Colour and Clarity of prepared medium

Purple coloured, clear solution without any precipitate

#### Reaction

Reaction of 2.8% w/v aqueous solution at 25°C. pH : 7.2±0.2

#### pH

7.00-7.40

#### Cultural Response

M1463: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Acid	Gas
<b>Cultural Response</b> <i>Alcaligenes faecalis</i> ATCC 8750	50-100	fair-good	negative reaction, no colour change	negative reaction

<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	positive reaction, yellow colour	positive reaction
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	good-luxuriant	positive reaction, yellow colour	positive reaction
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	good-luxuriant	positive reaction, yellow colour	positive reaction
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	good-luxuriant	positive reaction, yellow colour	positive reaction

## Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

## Reference

1. Eaton A. D., Clesceri L. S. and Greenberg A. E., (Eds), 1998, Standard Methods for the Examination of Water and Waste water, 20th Ed, APHA, Washington, D.C.
2. Corry J. E. L., Curtis G. D. W., and Baird R. M., Culture Media for Food Microbiology, Vol. 34, Progress in Industrial Microbiology, 1995, Elsevier, Amsterdam
3. Deutsche Einheitsverfahren zur Wasser- Abwasser- und Schalmuntersuchung. VCH Verlagsgesellschaft, D-6940, Weinheim.
4. Verordnung über Trinkwasser und über Wasser für Lebensmittelbetriebe vom 12. Dezember, 1990, Bundesgesetzbl., Teil I;2613-2629.1990.

Revision : 2 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.