



Antibiotic Assay Medium No. 36

M1666

Antibiotic Assay Medium No. 36 is used as a general purpose medium for cultivation of a wide variety of fastidious microorganisms.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	15.000
Papaic digest of soyabean meal	5.000
Sodium chloride	5.000
Agar	15.000
Final pH (at 25°C)	7.3±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 40.0 grams in 1000 ml purified / distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. If desired aseptically add 5% v/v defibrinated blood in previously cooled medium at 45 - 50°C. Mix well before pouring.

Principle And Interpretation

The composition of this medium is in accordance to CFR (1). This medium is recommended for sterility testing (2) Antibiotic Assay Medium No. 36 is widely employed as seed agar for agar diffusion assay for antibiotic bleomycin. The test organism *Mycobacterium smegmatis* is also maintained in this medium. This medium is employed for cultivation and isolation of fastidious or nonfastidious microorganisms. This medium is also used as maintenance medium of *Pseudomonas aeruginosa* for plate assay of ticarcillin. The medium like the conventional medium is used for a multitude of purposes including maintenance of stock cultures, plate counting, isolation of microorganisms from a variety of specimen types and a base for media containing blood (3,4).

Casein enzymic hydrolysate and papiac digest of soyabean gives the essential nutrients for maintenance and growth of the test organisms used. Osmotic balance is maintained by sodium chloride. Agar provides excellent medium for antibiotic diffusion and gives well defined zones of inhibition. Freshly prepared plates should be used for antibiotic assays. Test organisms are inoculated in sterile seed agar pre-cooled to 40-45°C and spread evenly over the surface of solidified base agar. All conditions in the microbiological assay must be controlled carefully. The use of standard culture media in the test is one of the important step for the good results.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Basal Medium : Light yellow coloured clear to slightly opalescent gel. After addition of 5-7% w/v sterile defibrinated blood : Cherry red coloured opaque gel forms in Petri plates

Reaction

Reaction of 4.0% w/v aqueous solution at 25°C (after sterilization). pH : 7.3±0.2

pH

7.10-7.50

Cultural Response

M1666: Cultural characteristics observed with added 5-7% w/v sterile defibrinated blood after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Growth w/ blood	Recovery w/ blood
Cultural Response					
Growth at 30-35°C for < = 3 days					
<i>Bacillus subtilis</i> ATCC 6633	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Staphylococcus aureus</i> ATCC25923	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Staphylococcus aureus</i> ATCC 6538	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Escherichia coli</i> ATCC 8739	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Escherichia coli</i> NCTC 9002	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Pseudomonas aeruginosa</i> ATCC 9027	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Salmonella Abony</i> NCTC 6017	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Micrococcus luteus</i> ATCC 9341	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Streptococcus pneumoniae</i> ATCC 6305	50-100	luxuriant	>=70%	luxuriant	>=70%
Growth at 20-25°C for < = 5 days					
* <i>Aspergillus brasiliensis</i> ATCC 16404	50-100	luxuriant		luxuriant	
<i>Candida albicans</i> ATCC 2091	50-100	luxuriant	>=70%	luxuriant	>=70%
<i>Candida albicans</i> ATCC 10231	50-100	luxuriant	>=70%	luxuriant	>=70%

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. Tests and Methods of Assay of Antibiotics and Antibiotic containing Drugs, FDA, CFR, 1983 Title 21, Part 436, Subpart D, Washington, D.C.: U.S. Government Printing Office, paragraphs 436, 100-436, 106, p. 242-259, (April 1).
2. Wright and Welch, 1959-60, Antibiotic Ann., 61.
3. MacFaddin 1985, Media for isolation-cultivation-identification-maintenance medical bacteria Vol, I, Williams, & Wilkins, Baltimore, MD
4. Forbes BA, Sahn DF, Weissfeld AS, 2002, Bailey and Scott's Diagnostic Microbiology, 11th ed.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.