



Lead Acetate Agar

M180

Lead Acetate Agar is recommended for the detection of hydrogen sulphide producing enteric bacteria.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	15.000
Proteose peptone	5.000
Dextrose	1.000
Lead acetate	0.200
Sodium thiosulphate	0.080
Agar	15.000
Final pH (at 25°C)	6.6±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 36.28 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Dispense into test tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Allow the tubes to cool in a slanted position to obtain slants with generous butts. Inoculate pure culture by surface streaking the slant and stabbing the butt.

Principle And Interpretation

Salmonella, *Shigella*, *Yersinia* species and certain strains of *Escherichia coli* cause severe gastroenteritis and life-threatening systemic illness in human (1, 2). Of these, *Salmonella* Typhi can be differentiated due to their ability to form hydrogen sulphide (3). Lead Acetate Agar is the modification of the original formulation of Spray (4). This medium was successfully used to study hydrogen sulphide production (4, 5). Lead Acetate Agar can also be used to differentiate between *Salmonella* Paratyphi A and *Salmonella* Paratyphi B (6). The latter produces hydrogen sulphide, observed as browning of the medium, within 18-24 hours, whereas the former fails to produce hydrogen sulphide.

Peptic digest of animal tissue, proteose peptone and dextrose provide all the essential nutrients for the growth of bacteria. Bacteria capable of using sulphur from sodium thiosulphate in their metabolic activities produce hydrogen sulphide. Lead acetate acts as an indicator of hydrogen sulphide production observed as browning of the medium. Dextrose is the fermentable carbohydrate source. Production of gas from dextrose is indicated by the presence of bubbles in the butt.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Medium amber coloured clear to slightly opalescent gel forms in tubes as slants

Reaction

Reaction of 3.63% w/v aqueous solution at 25°C. pH : 6.6±0.2

pH

6.40-6.80

Cultural Response

M180: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Gas Production	H ₂ S production
Cultural Response <i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	positive reaction	negative reaction

<i>Enterobacter aerogenes</i> ATCC 13048	50-100	luxuriant	positive reaction	negative reaction
<i>Salmonella Paratyphi A</i> ATCC 9150	50-100	luxuriant	negative reaction	negative reaction
<i>Salmonella Paratyphi B</i> ATCC 8759	50-100	luxuriant	negative reaction	positive reaction, browning of the medium
<i>Salmonella Typhi</i> ATCC 6539	50-100	luxuriant	variable reaction	positive reaction, browning of the medium
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	luxuriant	negative reaction	positive reaction, browning of the medium
<i>Shigella dysenteriae</i> ATCC 13313	50-100	luxuriant	negative reaction	negative reaction
<i>Shigella flexneri</i> ATCC 12022	50-100	luxuriant	negative reaction	negative reaction

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. Balows A., Hausler W. J. Jr., Hermann K. L., Isenberg H. D., Shadomy H. J., (Eds.), Manual of Clinical Microbiology, 5th Ed., ASM, Washington, D.C.
2. Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
3. Orłowski, 1897, Dissert., St. Petersburg.
4. Spray R. S., 1936, J. Bacteriol., 32:135.
5. Morrison L. E. and Tanner F. W., 1922, J. Bacteriol., 7:343.
6. Jordan E. O. and Victorson R., 1917, J. Infect. Dis., 21:554.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.