



Mueller Hinton Agar, 2% Glucose with Methylene blue

M1825

Mueller Hinton Agar, 2% Glucose with Methylene blue is recommended for testing performing Antifungal Disk Diffusion Susceptibility of yeasts.

Composition**

Ingredients	Gms / Litre
Beef infusion from	300.000
Casein Acid Hydrolysate	17.500
Starch	1.500
Glucose	20.000
Methylene blue	0.0005
Agar	17.000
Final pH (at 25°C)	7.3±0.1

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 58 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15lbs pressure (121°C) for 15 minutes. Mix well before pouring.

The performance of this batch has been tested and standardized as per the current CLSI (formerly NCCLS) document M44-A2 in Method for Antifungal Disk Diffusion susceptibility Testing of yeasts

Principle And Interpretation

The Mueller Hinton formulation was originally developed as a simple, transparent agar medium for the cultivation of pathogenic species (1). Mueller Hinton Agar, Modified (as per CLSI for antifungal) is recommended for the diffusion of antifungal agents impregnated on paper disc through an agar gel as described in CLSI Approved Standard (2).

When supplemented with glucose to a final concentration of 2%, it provides for suitable fungal growth. The addition of methylene blue to a final concentration of 5µg/ml enhances zone edge definition.

Kirby-Bauer et al recommended Mueller Hinton Agar for performing antibiotic susceptibility tests using a single disc of high concentration (4). WHO Committee on Standardization of Susceptibility Testing has accepted Mueller Hinton Agar for determining the susceptibility of microorganisms because of its reproducibility (3). Mueller Hinton Agar with 5% sheep blood and Mueller Hinton Agar with Haemoglobin have been recommended for antimicrobial susceptibility testing of *Streptococcus pneumoniae* and *Haemophilus influenzae*. Similarly Mueller Hinton Agar, Modified (as per CLSI for antifungal) is recommended for antifungal susceptibility testing of discs.

Beef infusion and casein acid hydrolysate provide nitrogenous compounds, carbon, sulphur and other essential nutrients. Starch acts as a protective colloid against toxic substances present in the medium. Starch hydrolysis yields dextrose, which serves as a source of energy. Glucose serves as an energy source for fungal cultures while Methylene blue enhances zone edge definition.

Technique:

Preparation of Inoculum:

1. Inoculum is prepared by picking five distinct colonies of approximately 1mm from 24 hours old culture grown on Sabouraud Dextrose Agar (M063) and incubated at $35 \pm 2^\circ\text{C}$. Colonies are suspended in 5ml of sterile 0.85% Saline.
2. Vortex the resulting suspension and adjust the turbidity to yield $1 \times 10^6 - 5 \times 10^6$ cells /ml (i.e. 0.5 McFarland standard).

Test Procedure:

1. Prepare plates with Mueller Hinton Agar, Modified (as per CLSI for antifungal) for carrying out susceptibility of antifungal discs. The medium in the plates should be sterile and have a depth of about 4 mm.

- Dip a sterile non-toxic cotton swab on a wooden applicator into the standardized inoculum (turbidity so adjusted, as to obtain semi confluent growth on the Petri plate) and rotate the soaked swab firmly against the upper inside wall of the tube to express excess fluid. Streak the entire agar surface of the plate with the swab three times, turning the plate at 60° angle between each streaking. Allow the inoculum to dry for 5 - 15 minutes with lid in place.
- Apply the discs using aseptic technique. Deposit the discs with centers at least 24 mm apart. (Not more than 12 discs should be placed on a 150-mm plate or not more than 5 discs on a 100-mm plate)
- Invert the plates and place in an incubator set to 35 ± 2°C within 15 minutes after the discs are applied.
- Examine each plate after 20 - 24 hours of incubation. If plate was satisfactorily streaked the resulting zones of inhibition will be uniformly circular and there will be a semi-confluent lawn of growth. Read at 48 hours only when insufficient growth is observed after 24 hours incubation.

Quality Control

Appearance

Light yellow to yellow may have slight blue tinge homogeneous free flowing powder

Gelling

Firm, comparable with 1.7% agar gel.

Colour and Clarity of prepared medium

amber coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 5.8% w/v aqueous solution at 25°C. pH : 7.3±0.1

pH

7.20-7.40

Cultural response

A luxuriant growth of test organisms was observed on Mueller Hinton Agar, Modified (as per CLSI for antifungal) in 24-48 hours at 33-37°C along with inhibition zones with respective antibiotic concentrations

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery	Amphotericin-B AP(100units)	Amphotericin-B AP(20 mcg)	Amphotericin-B AP(50 mcg)
Cultural response						
<i>Candida albicans</i> ATCC 90028	50-100	Luxuriant	≥70%	10 -17 mm	10 -15 mm	31 -42 mm
<i>Candida parapsilosis</i> ATCC 22019	50-100	luxuriant	≥70%	11 -20 mm	10 -17 mm	28 -37 mm
<i>Candida tropicalis</i> ATCC 750	50-100	luxuriant	≥70%	8 -12 mm	8 -10 mm	13 -17 mm
<i>Candida krusei</i> ATCC 6258		luxuriant	≥70%	9 -14 mm	8 -12 mm	16 -25 mm
<i>Candida albicans</i> ATCC 10231	50-100	luxuriant	≥70%	10 -18 mm	10 -16 mm	30 -40 mm
<i>Saccharomyces cerevisiae</i> ATCC9763	50-100	luxuriant	≥70%	11 -18 mm	8 -12 mm	29 -38 mm

Storage and Shelf Life

Store dehydrated powder below 30°C and prepared medium at 2-8°C. Use before expiry date on the label.

Reference

- Mueller J. H. and Hinton J., 1941, Proc. Soc. Exp. Biol. Med., 48:330.
- Method for Antifungal Disk Diffusion Susceptibility Testing of yeasts; Approved Guideline Second Edition M44-A2 Vol.24 No.17.
- Present Status and Future Work, WHO Sponsored collaborative study, Chicago, Oct. 1967.
- Bauer A. W., Kirby W. M., Sherris J. L. and Turck M., 1966, Am. J. Clin. Pathol., 45:493.

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.