



HiCrome™ Bifidobacterium Agar

M1960

Intended use

Recommended for the differentiation of *Bifidobacterium* and *Lactobacillus* species from clinical and non-clinical samples.

Composition**

Ingredients	Gms / Litre
Peptone special	23.000
Sodium chloride	5.000
M-Protein powder #	5.000
Chromogenic mixture	10.480
Agar	16.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

- Equivalent to Milk powder

Directions

Suspend 59.48 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.

Principle And Interpretation

The genus *Bifidobacterium* is the third most numerous bacterial populations found in the human intestine after *Bacteroides* and *Eubacterium*. It is an anaerobic bacteria that makes up the gut microbial flora. It resides in the colon and have health benefits for their hosts. Bifidobacteria are also associated with lower incidences of allergies (3,4). Bifidobacterium Agar is used for the cultivation and maintenance of *Bifidobacterium* species (2).

Peptone special provides nitrogenous and carbonaceous compounds, long chain amino acids, vitamins and other essential growth nutrients. Sodium chloride maintains osmotic balance. M-protein powder aids in detecting casein hydrolysis activity which is exhibited by *Bifidobacterium breve*. A halo zone is observed around the colony in case of casein hydrolysis. The indicator system in the chromogenic mixture helps in distinguishing between *Lactobacillus* and *Bifidobacterium* species. *Lactobacillus* species usually produce green colonies with opaque zone. *Bifidobacterium infantis* produces dark blue to bluish green colonies. Agar serves as a solidifying agent.

Type of specimen

Clinical samples - Faeces ; Food and dairy samples.

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (5,6).

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,7,8).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Warning and Precautions :

In Vitro diagnostic Use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations :

1. Due to variable nutritional requirements, some strains may show poor growth on this medium.
2. Slight colour variation may be observed depending upon the utilization of the substrate by the organism.
3. *Bifidobacterium* species are strict anaerobes, hence condition must be appropriately maintained.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.6% Agar gel

Colour and Clarity of prepared medium

Reddish orange coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 5.95% w/v aqueous solution at 25°C. pH :7.2±0.2

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 48 hours under anaerobic conditions.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Bifidobacterium infantis</i> ATCC 25962	50-100	good-luxuriant	≥50%	Dark blue-bluish green
<i>Bifidobacterium breve</i> ATCC 50-100 15698	50-100	good-luxuriant	≥50%	Red-Pink with halo zone
<i>Lactobacillus plantarum</i> ATCC 8014	50-100	good-luxuriant	≥50%	Green colonies w/ hazy background
<i>Lactobacillus fermentum</i> ATCC 9338	50-100	good-luxuriant	≥50%	Pink without halo zone

Storage and Shelf Life

Store the dehydrated and the prepared medium in a tightly closed container at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (5,6).


Reference

1. American Public Health Association, Standard Methods for the Examination of Dairy Products, 1978, 14th Ed., Washington D.C.
2. Atlas R. M. 2004, 3rd Edi. Handbook of Microbiological Media, Parks, L. C. (Ed.), CRC Press, Boca Raton.
3. Björkstén B., Sepp E., Julge K., Voor T., and Mikelsaar M., 2001, J. Allergy Clin. Microbiol., Volume 108, Issue 4, 516-520.
4. Guarner F., and Malagelada J. R., 2003, The Lancet, Vol. 361, Issue 9356, 8 February 2003, 512-519.
5. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
6. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
7. Salfinger Y., and Tortorello M.L. Fifth (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 5th Ed., American Public Health Association, Washington, D.C.
8. Wehr H. M. and Frank J. H., 2004, Standard Methods for the Microbiological Examination of Dairy Products, 17th Ed., APHA Inc., Washington, D.C.

IVD



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,
23 Vadhani Industrial Estate,
LBS Marg, Mumbai-86, MS, India



CE Partner 4U ,Esdoornlaan 13, 3951
DB Maarn The Netherlands,
www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.