



APT Agar

M226

APT Agar is recommended for the cultivation of heterofermentative lactic acid bacteria and other organisms requiring high thiamine content.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	12.500
Yeast extract	7.500
Dextrose	10.000
Sodium citrate	5.000
Sodium chloride	5.000
Dipotassium phosphate	5.000
Magnesium sulphate	0.800
Manganese chloride	0.140
Ferrous sulphate	0.040
Polysorbate 80	0.200
Thiamine hydrochloride	0.001
Agar	15.000
Final pH (at 25°C)	6.7±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 61.18 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Dispense as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. AVOID EXCESSIVE HEATING.

Principle And Interpretation

APT (All purpose Tween 80) Agar is formulated as per Evans and Niven (1) for cultivation and maintenance of Lactobacilli. This medium is also used in the microbiological assay of thiamine. Lactobacillus forms a major part of lactic acid bacteria group which are abundant in nature. They convert lactose and other sugars to lactic acid and therefore are named as *Lactobacillus*. They are responsible for spoilage of foods like meat, dairy etc. However APT Agar can also be used for cultivation of heterofermentative lactic acid bacteria requiring high thiamine content (2). APT Agar is also used as a maintenance medium since it preserves the viability and sensitivity of *Lactobacillus viridescens* ATCC 12706.

APT Agar contains casein enzymic hydrolysate, which acts as a source of carbon, nitrogen, vitamins and minerals. Yeast extract provides vitamin and B-complex nutrients, which is required for the growth of bacteria. Dextrose is the carbohydrate source. Manganese chloride, magnesium sulphate and ferrous sulphate provide ions used in replication by lactobacilli. Polysorbate 80 is a source of fatty acids required by lactobacilli.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 6.12% w/v aqueous solution at 25°C. pH : 6.7±0.2

pH

6.50-6.90

Cultural Response

M226: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Recovery
Cultural Response			
<i>Lactobacillus acidophilus</i> ATCC 4356	50-100	good-luxuriant	>=50%
<i>Lactobacillus viridescens</i> ATCC 12706	50-100	good-luxuriant	>=50%
<i>Leuconostoc mesenteroides</i> ATCC 12291	50-100	good-luxuriant	>=50%
<i>Lactobacillus casei</i> ATCC 9595	50-100	good-luxuriant	>=50%
<i>Lactobacillus plantarum</i> ATCC 8014	50-100	good-luxuriant	>=50%

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on label.

Reference

1. Evans and Niven, 1951, J. Bact., 62:599.
2. Downes F. P. and Ito K. (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th ed., APHA, Washington D.C.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.