



Azotobacter Agar (Glucose)

M371

Azotobacter Agar (Glucose) is recommended for isolation and cultivation of Glucose positive *Azotobacter* species from soil.

Composition**

Ingredients	Gms / Litre
Dipotassium phosphate	1.000
Magnesium sulphate	0.200
Sodium chloride	0.200
Ferrous sulphate	0.005
Soil extract	5.000
Glucose	10.000
Agar	15.000
Final pH (at 25°C)	7.6±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 31.4 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. If slight precipitate occurs after autoclaving, distribute it evenly before pouring into sterile Petri plates.

Principle And Interpretation

Bacteria of the family *Azotobacteraceae* constitute the majority of heterotrophic free-living nitrogen fixing bacteria (1). *Azotobacter* is a genus of free-living diazotrophic bacteria which have the highest metabolic rate compared to any other microorganisms. *Azotobacters* have generated a good deal of interest in the scientific community because of their unique mode of metabolism, by which they can fix nitrogen aerobically. Azotobacter Agar (Glucose) is used for isolation and cultivation of glucose positive *Azotobacter* species from soil (2). It is also useful for maintenance of *Azotobacter* species by adding extra 1% glucose to the medium as specified by the American Type Culture Collection (3).

Quality Control

Appearance

Off white to beige homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel with slight precipitate forms in Petri plates

Reaction

Reaction of 3.14% w/v aqueous solution at 25°C. pH : 7.6±0.2

pH

7.40-7.80

Cultural Response

M371: Cultural characteristics observed after an incubation at 25-30°C for 24-48 hours or longer.

Organism

Growth

Cultural Response

Azotobacter beijerinckii good-luxuriant

ATCC 12981

Azotobacter nigricans ATCC good-luxuriant

35009

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label

Reference

1. Subba Rao N. S., 1977, Soil Microorganisms and Plant Growth, Oxford and IBH Publishing Co., New Delhi.
2. Pelczar M. Jr., 1957, Manual of Microbiological Methods.
3. ATCC Catalogue of Bacteria and Bacteriophages, 1992, 18th Ed, American Type Culture Collection, Rockville, MD.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
