



Halophilic Broth

M591

Halophilic Broth is used for the isolation and cultivation of extremely halophilic bacteria.

Composition**

Ingredients	Gms / Litre
Casein acid hydrolysate	10.000
Yeast extract	10.000
Proteose peptone	5.000
Trisodium citrate	3.000
Potassium chloride	2.000
Magnesium sulphate	25.000
Sodium chloride	250.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 30.5 grams in 100 ml distilled water. Heat if necessary to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

Halophilic media are formulated for isolation and cultivation of extreme halophilic species of *Halobacterium* and *Halococcus* from foods (1, 2). For optimum growth they require high salt concentration of about 20 - 30%. In general, the requirement for salt by halophilic microorganisms is not an exclusive need for NaCl since many species require low levels of K⁺, Mg⁺⁺ and other ions in addition to NaCl (3, 4). The level of salt required by microorganism varies greatly. Therefore the microbial types associated with a particular salted food depend on the concentration and type of salt and food. The most recent classifications of halophilic microorganisms are based on the level of salt required (2, 3). These bacteria can cause pink discoloration on the outer surface accompanied by putrefaction and decomposition of fish, bacon and hides preserved in sea salts.

Halophilic Broth contains casein acid hydrolysate; proteose peptone and yeast extract which provide all the necessary nutrients, mainly nitrogenous and vitamins to the halophilic bacteria. Trisodium citrate is added to avoid the losses (2). Magnesium sulphate, sodium chloride and potassium chloride are essential ions required for the growth of extreme halophiles.

10 gm sample is added to 90 ml Halophilic Broth and incubated at 35°C for upto 12 days. The organisms are then isolated onto Halophilic Agar (M590) from this enriched culture.

Quality Control

Appearance

Off-white to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Amber coloured, slightly hazy solution with heavy precipitate at the bottom in tubes.

Reaction

Reaction of 30.5% w/v solution at 25°C. pH : 7.2±0.2

pH

7.00-7.40

Cultural Response

M591: Cultural characteristics observed after an incubation at 35-37°C for 12 days.

Organism

Growth

Halobacterium salinarium luxuriant
ATCC 33171
Halococcus morrhuae ATCC luxuriant
17082

Storage and Shelf Life

Store below 30°C in tightly closed container and prepared medium at 2-8°C. Use before expiry period on the label.

Reference

1. Dundas I.E., 1977, Advances In Microbiology and Physiology, Rose H. and Tempest D.W. (Eds.), A.P. London.
2. Gibbons N.E., 1969, Methods In Microbiology, Vol. 3B, Norris J.R., and Ribbons D.W. (Eds.), A.P., New York, pp.169-183.
3. Kushner D. J., (Eds.), 1978, D. J. Kushner, pg 317, Academic Press, London, England
4. MacLeod R. A., 1965, Bacteriol., Rev., 29:9

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.