



Casman Broth Base

M766

Casman Broth Base with blood is used for isolation of fastidious microorganisms from clinical specimens under reduced oxygen tension.

Composition**

| Ingredients | Gms / Litre |
|-----------------------------|-------------|
| Proteose peptone | 10.000 |
| Tryptose | 10.000 |
| Beef extract | 3.000 |
| Dextrose | 0.500 |
| Corn starch | 1.000 |
| Sodium chloride | 5.000 |
| Nicotinamide | 0.050 |
| p-Amino benzoic acid (PABA) | 0.050 |
| Final pH (at 25°C) | 7.2±0.2 |

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 29.6 grams in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 50°C and aseptically add 0.15% v/v sterile waterlysed blood (water:blood :: 3:1) of 5% sterile blood. Alternatively add 5% partially lysed blood. Mix well and dispense as desired.

Principle And Interpretation

Fastidious microorganisms such as *Haemophilus* and *Neisseria* require the addition of X and V- growth factors for in vitro cultivation (1). Casman (1, 2, 3) described a blood-enriched medium for cultivation of *Haemophilus* and gonococci (1). The medium was developed to replace the previously described formulations that required time-consuming preparations using fresh and heated blood and meat infusion to supply the essential nutrients for growth of these fastidious organisms (2, 3). Blood supplies factor-X (hemin) and factor-V (Nicotinamide Adenine Dinucleotide), which is required for growth of *Haemophilus influenzae*. Sheep blood lacks factor-V due to NADase, an enzyme that destroys factor- V (4). Horse and rabbit blood supplies both the factor X and factor V, and are relatively free of NADase activity, therefore it is preferred over sheep blood. Nicotinamide is added to medium to inhibit nucleotidase of erythrocytes that may destroy factor V.

Proteose peptone, tryptose and beef extract provide amino acids and other complex nitrogenous nutrients. Dextrose improves growth of pathogenic cocci. Corn starch prevents fatty acids from inhibiting the growth of *Neisseria gonorrhoeae*, without interfering with haemolytic reaction. Corn starch also neutralizes the inhibitory action of dextrose. Inoculate the medium as soon as the specimen arrives at the laboratory.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Basal medium: Yellow coloured clear to slightly opalescent solution. After addition of blood: Cherry red coloured opalescent solution in tubes

Reaction

Reaction of 2.96% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH

7.00-7.40

Cultural Response

M766: Cultural characteristics observed with added water-lysed blood, after an incubation at 35-37°C for 40-48 hours.

| Organism | Inoculum (CFU) | Growth |
|---|----------------|----------------|
| <i>Haemophilus influenzae</i> ATCC 35056 | 50-100 | good-luxuriant |
| <i>Neisseria meningitidis</i> ATCC 13090 | 50-100 | luxuriant |
| <i>Streptococcus mitis</i> ATCC 9895 | 50-100 | luxuriant |
| <i>Streptococcus pneumoniae</i> ATCC 6303 | 50-100 | luxuriant |
| <i>Streptococcus pyogenes</i> ATCC 19615 | 50-100 | luxuriant |

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. Casman, 1947, Am. J. Clin. Pathol., 17:281.
2. Casman, 1942, J. Bact., 43:33.
3. Casman, 1947, J. Bact., 53:561.
4. Krunveide and Kuttner, 1938, J. Exp. Med., 67:429.

Revision : 1 / 2011



Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.