



## Antibiotics Assay Medium No. 38

M799

Antibiotic Assay Medium No. 38 is used for the microbiological assay of Ticarcillin, using *Pseudomonas aeruginosa* as the test organism.

### Composition\*\*

| Ingredients                              | Gms / Litre |
|--|-------------|
| Peptic digest of animal tissue (Peptone) | 15.000      |
| Papaic digest of soyabean meal           | 5.000       |
| Sodium chloride                          | 4.000       |
| Sodium sulphite                          | 0.200       |
| L-Cystine                                | 0.700       |
| Dextrose                                 | 5.500       |
| Agar                                     | 15.000      |
| Final pH ( at 25°C)                      | 7.0±0.2     |

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 45.4 grams in 1000 ml purified/distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

### Principle And Interpretation

This medium follows the specification of CFR (1) and is routinely employed for agar diffusion assay of Ticarcillin using Gram negative test organisms specially *Pseudomonas aeruginosa*. This medium is used as both base agar and seed agar for assay of Ticarcillin.

Peptic digest of animal tissue and papaic digest of soyabean meal provides essential nutrients and growth factors for the growth of test organisms. Dextrose serves as carbon source. Sodium chloride maintains the osmotic equilibrium. L-cystine and sodium sulphite are sulphur providers that aids assimilation of sulphur during microbial growth. L-cystine also acts as growth stimulator and enrich the medium with amino acid source for promoting the growth. The high nutritional content along with high sulfur (cystine and sodium sulphite) content improves growth with chromogenicity of test organism *Pseudomonas*.

Freshly prepared plates should be used for antibiotic assays. Test organisms are inoculated in sterile seed agar pre-cooled to 40-45°C and spread evenly over the surface of solidified base agar.

### Quality Control

#### Appearance

Cream to yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 1.5% Agar gel

#### Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in Petri plates.

#### Reaction

Reaction of 4.54% w/v aqueous solution at 25°C. pH : 7.0±0.2

#### pH

6.80-7.20

#### Cultural Response

M799: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours .

| Organism | Inoculum (CFU) | Growth | Recovery | Antibiotics assayed |
|----------|----------------|--------|----------|---------------------|
|----------|----------------|--------|----------|---------------------|

*Pseudomonas aeruginosa* 50-100 luxuriant  $\geq 70\%$  Ticarcillin  
ATCC 29336

### Storage and Shelf Life

Store below 30°C in tightly closed container and use freshly prepared medium. Use before expiry date on the label

### Reference

1. Tests and Methods of Assay of Antibiotics and Antibiotic containing Drugs, FDA, CFR, 1983 Title 21, Part 436, Subpart D, Washington, D.C.: U.S. Government Printing Office, paragraphs 436, 100-436, 106, p. 242-259, (April)

Revision : 2 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.