



## Nitrofurantoin Broth Base

M857

Nitrofurantoin Broth Base is used for the selective enrichment and isolation of *Pseudomonas* species.

### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	7.500
Casein enzymic hydrolysate	7.500
Sodium chloride	5.000
Final pH ( at 25°C)	7.2±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 20 grams in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to room temperature and aseptically add 50 ml sterile 0.2% nitrofurantoin solution. Mix well and dispense in tubes or flasks as desired. Sterile nitrofurantoin solution (0.2%) is prepared by dissolving 1 gm Nitrofurantoin in 500 ml polyethylene glycol 300.

Note: Autosterilization takes place in 3 months. This solution can be stored for 6 months or longer.

### Principle And Interpretation

Selective and differentiating media consisting of simple chemical components have been developed, both in solid and in liquid form, for culturing *Pseudomonas aeruginosa*. Nitrofurantoin, in the form of Macrochantin, has been shown to be active against most strains of *Escherichia coli*, *Staphylococcus aureus* and *Enterococcus faecalis* both in vitro and in clinical infections. Nitrofurantoin is not active against most strains of *Proteus* species or *Serratia* species. It has no activity against *Pseudomonas* species (1). Therefore nitrofurantoin incorporated in medium can be used as a selective medium for culturing of *Pseudomonas* species.

Casein enzymic hydrolysate and peptic digest of animal tissue provide the essential nutrients especially nitrogenous sources. Nitrofurantoin, 1-[5-nitrofururylidene) amino] hydantoin, is a synthetic antibacterial agent which is effective against most common gram-negative and gram-positive urinary tract pathogenic bacteria (2).

### Quality Control

#### Appearance

Cream to yellow homogeneous free flowing powder

#### Colour and Clarity of Prepared medium

With added nitrofurantoin : Fluorescent yellow coloured clear solution without any precipitate

#### Reaction

Reaction of 2.0% w/v aqueous solution at 25°C. pH : 7.2±0.2

#### pH

7.00-7.40

#### Cultural Response

M857: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth
<i>Escherichia coli</i> ATCC 25922	≥10 <sup>3</sup>	inhibited
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	good-luxuriant
<i>Staphylococcus aureus</i> ATCC 25923	≥10 <sup>3</sup>	inhibited

## Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

## Reference

1. Clinical and Laboratory Standards Institute, 2006, Performance standards for antimicrobial susceptibility testing. Approved standard M100-S15, Vol. 25, CLSI, Villanova, Pa.
2. Chamberlain R. E., 1976, Chemotherapeutic properties of prominent nitrofurans, J. Antimicrob. Chemother. 2:325336

Revision : 2 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.