



## Motility Test Medium (Edwards and Ewing)

M930

Motility Test Medium (Edwards and Ewing) is used for testing motility of enteric bacteria.

### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	10.000
Beef extract	3.000
Sodium chloride	5.000
Agar	4.000
Final pH ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 22 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Dispense 8 ml amounts in test tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool the tubed medium in an upright position.

### Principle And Interpretation

Bacterial motility can be observed directly on microscopic slide or it can be visualized on motility media having agar concentration of 0.4% or less (1). Use of such semisolid media to observe or detect motility was reported by Tittsler and Sandholzer (2). Motility Test Medium is the modification of the original formulation as per Edwards and Ewing and is used for testing motility of *Enterobacteriaceae* (3). Motility can be visualized as a diffused zone of growth flaring out from the line of inoculation (1,4,5).

Peptic digest of animal tissue, beef extract serve as sources of essential growth nutrients required for bacterial metabolism. Sodium chloride maintains the osmotic equilibrium of the medium. Small amount of agar helps to create a semisolid medium.

Bacterial motility can be observed directly by examination of the tubes following incubation. Inoculation is done by stabbing through the centre of the medium. Incubate at appropriate temperature for 18 to 40 hours. Non-motile organisms grow only along the line of inoculation whereas motile organisms grow away from the line of inoculation or may show growth even throughout the medium. All weak or equivocal motility results should be confirmed by flagellum stain or by direct wet microscopy (hanging drop) (6, 7). To enhance the visibility of bacterial growth 2,3,5 Triphenyl Tetrazolium Chloride (TTC) (FD057) may be added. Tetrazolium salts are colourless but are converted into insoluble formazan, a red coloured complex by the reducing properties of growing bacteria. In Motility Test Medium containing tetrazolium, the development of this red colour helps to trace the spread of bacteria from the inoculation line. The motility of *Listeria monocytogenes* is frequently best observed in medium without TTC.

### Quality Control

#### Appearance

Cream to yellow homogeneous free flowing powder

#### Gelling

Semisolid, comparable with 0.4% Agar gel.

#### Colour and Clarity of prepared medium

Light yellow coloured clear to slightly opalescent gel forms in tubes as butts

#### Reaction

Reaction of 2.2% w/v aqueous solution at 25°C. pH : 7.4±0.2

#### pH

7.20-7.60

#### Cultural Response

M930: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours .

Organism	Inoculum (CFU)	Growth	Motility
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	positive, growth away from stabline causing turbidity
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	luxuriant	positive, growth away from stabline causing turbidity
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	luxuriant	negative, growth along the stabline, surrounding medium remains clear
<i>Salmonella Enteritidis</i> ATCC 13076	50-100	luxuriant	positive, growth away from stabline causing turbidity
<i>Staphylococcus aureus</i> ATCC 25923	50-100	luxuriant	negative, growth along the stabline, surrounding medium remains clear
<i>Vibrio cholerae</i> ATCC 15748	50-100	luxuriant	positive, growth away from stabline causing turbidity
<i>Vibrio parahaemolyticus</i> ATCC 17802	50-100	luxuriant	positive, growth away from stabline causing turbidity

## Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

## Reference

1. Koneman E. W., Allen S. D., Janda W. M., Schreckenberger P. C., Winn W. C. Jr., (Eds.), 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company.
2. Tittsler R. P. and Sandholzer L. A., 1936, J. Bacteriol., 31:575.
3. Edward P. R. and Ewing W. H. 1972, Identification of Enterobacteriaceae 3rd Ed., Minneapolis, Burgess.2.,
4. Howard B. J. and Other (Eds.), 1994, Clinical and Pathogenic Microbiology, The C. V. Mosby. Year Book, Inc.
5. Baron. E. J. and Finegold S. M. (Eds.), 1990, Bailey and Scott's `Diagnostic Microbiology, 8th ed., The C. V. Mosby. Co, St., Louis, Missouri.
6. DAmato R. F., and Tomfohrede K. M., 1981, J. Clin. Microbiol., 14 (3), 347-348.
7. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore.

Revision : 2 / 2015

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.