



## KF Streptococcus Agar Base w/ BCP

M1007

KF Streptococcus Agar Base w/ BCP is recommended for detection and enumeration of faecal Streptococci.

### Composition\*\*

Ingredients	Gms / Litre
Proteose peptone	10.000
Yeast extract	10.000
Sodium chloride	5.000
Sodium glycerophosphate	10.000
Maltose	20.000
Lactose	1.000
Sodium azide	0.400
Bromocresol purple	0.015
Agar	20.000
Final pH ( at 25°C)	7.2±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 76.41 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. DO NOT AUTOCLAVE. Overheating will lower the pH and render the medium less productive. Cool to 45-50°C and aseptically add 10 ml of 1% 2, 3, 5-Triphenyl Tetrazolium Chloride (TTC) (FD057) to sterile medium. Mix well and pour into sterile Petri plates.

### Principle And Interpretation

Streptococci are spherical, gram-positive bacteria and form a part of the normal commensal flora of the mouth, skin, intestine, upper respiratory tract of humans. Streptococci found in the faeces form the faecal Streptococci and constitute of Streptococci with group D Lancefield antigens. The types include *Streptococcus faecalis*, *Streptococcus faecium*, *Streptococcus bovis* and *Streptococcus duran*. They are low-grade pathogens and rarely cause disease. However, they may cause urinary tract infection in catheterized patients; mixed abdominal wound infections following gut surgery; and endocarditis on abnormal valves. Kenner - Faecal (KF) Medium were developed by Kenner et al (1, 2) for detecting Streptococci in water and food materials.

Proteose peptone along with yeast extract provide nitrogen, carbon, sulphur, amino acids, vitamins and trace ingredients to the faecal Streptococci. Lactose and maltose are the fermentable carbohydrates and therefore serve as energy sources. Sodium azide is a selective agent, which hampers the growth of gram-negative bacteria.

2,3,5-Triphenyl Tetrazolium Chloride is reduced to insoluble formazan by actively metabolizing cells, resulting in the formation of pink or red colonies. Bacteria resistant to azide, utilize lactose and / or maltose. The acidity so produced changes the colour of the indicator dyes to yellow. Bacterial cells reduce TTC to insoluble formazan, resulting in the formation of pink to red colonies.

Samples can be directly streaked or sterile membrane filters through which the water samples have been passed are aseptically placed on the media. After an incubation at 35-37°C for 24-48 hours, Enterococci appear as pink to red colonies. After this presumptive identification, further confirmatory tests should be carried out (3,4).

### Quality Control

#### Appearance

Cream to greyish yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 2.0% agar gel.

#### Colour and Clarity of prepared medium

Light purple coloured, clear to slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 7.64% w/v aqueous solution at 25°C. pH : 7.2±0.2

#### pH

7.00-7.40

#### Cultural Response

M1007: Cultural characteristics observed with added FD057, after an incubation at 35-37°C for 48-72 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<b>Cultural Response</b>				
<i>Enterobacter aerogenes</i> ATCC 13048	≥10 <sup>3</sup>	inhibited	0%	
<i>Enterococcus faecalis</i> ATCC 50-100 29212		good-luxuriant	≥50%	red-maroon
<i>Escherichia coli</i> ATCC 25922	≥10 <sup>3</sup>	inhibited	0%	

#### Storage and Shelf Life

Store below 30°C in tightly closed container and prepared medium at 2-8°C. Use before expiry period on the label.

#### Reference

1. Kenner B. A., Clark H. F. and Kabler P. W., 1960, Am. J. Public Health, 50:1553.
2. Kenner B. A., Clark H. F. and Kabler P. W., 1961, Appl. Microbiol., 9:15.
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore.
4. Facklam R. R. and Moody M. P., 1970, Appl. Microbiol., 20:245.

Revision : 1 / 2011



#### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.