



C.L.E.D. Agar Base w/o Indicator

M1146

Intended Use:

C.L.E.D. Agar w/o Indicator (with added Bromo Thymol Blue) is recommended for isolation, enumeration and presumptive identification of bacterial flora in the urinary tract.

Composition**

Ingredients	Gms / Litre
Peptone	4.000
Casein enzymic hydrolysate	4.000
Meat Extract B #	3.000
Lactose	10.000
L-Cystine	0.128
Agar	15.000
Final pH (at 25°C)	7.3±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 36.1 grams in 998 ml distilled water. Add rehydrated contents of 1 vial of Bromo Thymol Blue Supplement (FD091). Heat, to boiling, to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

Principle And Interpretation

On a solid medium, Sandy's reported that swarming of *Proteus* species could be controlled by restricting the electrolytes (8). Formerly swarming of *Proteus* was controlled by adding alcohol, surface-active agent, sodium azide, boric acid etc. to the medium (8). Later on Sandys medium was modified by Mackey and Sandy's (6), by replacing mannitol by lactose and sucrose and elevating concentration of agar and bromothymol blue. This formulation was further modified by the same authors and called C.L.E.D. (Cystine-Lactose-Electrolyte-Deficient) by deleting the sucrose and by including L-cystine for promoting the growth of cystine dependent dwarf coliform colony (7). This medium is recommended for use in urine bacteriology, promoting the growth of all urinary pathogens. C.L.E.D. Medium is also recommended for dipstick procedures and as dip inoculum transport medium for urine specimens (2,6,7).

Peptone, meat extract B and casein enzymic hydrolysate provides nitrogen and carbon source, long chain amino acids, vitamins and other essential growth nutrients. Lactose is the fermentable sugar. L-cystine supports the growth of dwarf coliform colony. Bromo thymol blue is the pH indicator which turns yellow at acidic pH.

Bacteriuria may be quantitated by inoculating the surface of an agar medium by proper dilution. Inoculate the medium immediately after urine collection. It can also be inoculated by calibrated loop or duplicate dilution pour plate methods (1,5). *Shigella* species may not grow on this medium. Initiation of antibiotic therapy, before collection sample, low urine pH (less than 5) etc. may result in low bacterial count from infected patients.

Type of specimen

Clinical:Urine

Specimen Collection and Handling

For clinical samples follow appropriate techniques for handling specimens as per established guidelines (3,4).

Warning and Precautions

In Vitro diagnostic use only. Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling clinical specimens. Safety guidelines may be referred in individual safety data sheets.

Limitations

- 1.This medium is recommended for urine infection. Low urine count may be a result of antibiotic therapy, low pH of urine.
- 2.Recovery depends on the urine count.
- 3.Inoculate the medium immediately after urine collection.
4. *Shigella* species may not grow on this medium.
- 5.For better results, the medium should not be incubated for more than 24 hours because if lactose fermenters predominate the entire medium may turn yellow masking the presence of non-lactose fermenters.

Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

With addition of Bromo Thymol Blue Supplement (FD091): Green coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 3.61% w/v aqueous solution at 25°C. pH : 7.3±0.2

pH

7.10-7.50

Cultural Response

Cultural characteristics observed with added Bromothymol Blue Supplement(FD091),after an incubation at 35-37°C for 18-24 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
Cultural Response <i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	≥70%	yellow, opaque, center slightly deeper yellow
<i>Enterococcus faecalis</i> ATCC 29212	50-100	good-luxuriant	≥70%	slight yellowish or greenish
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	good-luxuriant	≥70%	yellow to whitish blue
<i>Proteus vulgaris</i> ATCC 13315	50-100	good-luxuriant	≥70%	blue
<i>Staphylococcus aureus</i> ATCC 25923	50-100	good-luxuriant	≥70%	deep yellow
<i>Salmonella Typhi</i> ATCC 6539	50-100	good-luxuriant	≥70%	bluish

Storage and Shelf Life

Store below 30°C in a tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label. On opening, product should be properly stored dry,after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with clinical sample must be decontaminated and disposed of in accordance with current laboratory techniques (3,4).

Reference

1. Benner E. J., 1970, Appl. Microbiol., 19(3), 409
2. Dixon J. M. S. and Clark M. A., 1968, Conc. Med. Assoc. J., 99 (15)
3. Isenberg, H.D. Clinical Microbiology Procedures Handbook. 2nd Edition.
4. Jorgensen, J.H., Pfaller, M.A., Carroll, K.C., Funke, G., Landry, M.L., Richter, S.S and Warnock., D.W. (2015) Manual of Clinical Microbiology, 11th Edition. Vol. 1.
5. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore
6. Mackey and Sandys, 1965, Br. Med. J., 2:1286.
7. Mackey and Sandys, 1966, Br. Med. J., 1:1173.
8. Sandys, 1960, J. Med. Lab. Technol., 17:224.


Revision : 02 / 2016



In vitro diagnostic medical device



CE Marking



Storage temperature



Do not use if package is damaged



HiMedia Laboratories Pvt. Limited,
B /4-6 , MIDC, Dindori, Nashik MH

www.himedialabs.com



CE Partner 4U ,Esdoornlaan 13, 3951
DB Maarn The Netherlands,

www.cepartner4u.eu

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.