

## Milk Agar with Cetrimide (Twin Pack)

**M1273**

Milk Agar with Cetrimide is used for the detection and enumeration of *Pseudomonas aeruginosa* in swimming pool waters.

### Composition\*\*

Ingredients	Gms / Litre
Part A	-
Skim milk powder	133.330
Part B	-
Peptic digest of animal tissue	3.330
Sodium chloride	1.670
Yeast extract	1.000
Cetrimide	0.400
Agar	20.000
Final pH ( at 25°C )	7.3±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 26.4 grams of Part B in 250 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 20 minutes. Suspend 133.33 grams of Part A in 750 ml of distilled water and sterilize by autoclaving at 15 lbs pressure (121°C) for 5 minutes. After autoclaving mix Part A and B and pour into sterile Petri plates.

### Principle And Interpretation

Milk Agar was modified by Brown and Scott (1) for the confirmation of *Pseudomonas aeruginosa* in swimming pool waters. Swimming pool water is generally chlorinated potable water but it can also be from thermal springs or salt water. Microorganisms of concern are typically those from the body of the bathers, including the orifices. As *P.aeruginosa* can survive for longer time in water compared to other microorganisms, it is one of the major indicator organisms in the swimming pool. This organism is mainly responsible for ear and eye infection and is very likely to get disseminated in the swimming pool water due to constant contact of ears and eyes with the water.

Milk Agar with Cetrimide is formulated in accordance with ISO Committee under the specifications ISO 8360-1: 1988 for the detection and enumeration of *P.aeruginosa* from water (2). Strains of *P.aeruginosa* are identified by their pigment production i.e. pyocyanin. *P.aeruginosa* is the only species of *Pseudomonas* or gram-negative rod known to excrete pyocyanin.

*P.aeruginosa* hydrolyzes casein and produces a yellowish to green diffusible pigment on Milk Agar. For isolation, filter 200ml or less water of the swimming pool through sterile membrane filters. Place each membrane filter on M-PA Agar (M1121). Incubate the plates at 41.5±0.5°C for 72 hours. Typical *P.aeruginosa* colonies are 0.8-2.2 mm in diameter, flat in appearance with brownish to greenish centers. For confirmation, using Milk Agar w/ Cetrimide, make a single streak from an isolated colony on a Milk Agar w/ Cetrimide plate and incubate at 35-37°C for 24 hours. After incubation *P.aeruginosa* forms pigmented colonies.

Milk, skim milk powder, peptic digest of animal tissue and yeast extract provide all the necessary nutrients mainly nitrogenous for the multiplication of *P.aeruginosa*.

*P.aeruginosa* forms yellowish green colonies on this medium. Cetrimide acts as a quaternary ammonium, cationic detergent that causes release of nitrogen and phosphorus from bacterial cells other than *P.aeruginosa*.

### Quality Control

#### Appearance

Part A : White to cream homogeneous free flowing powder Part B : Cream to yellow homogeneous free flowing powder

**Gelling**

Firm, comparable with 2.0% Agar gel.

**Colour and Clarity of prepared medium**

Light amber coloured opalescent gel forms in Petri plates

**Reaction**

Reaction of 2.64% w/v aqueous solution of Part B at 25°C. pH : 7.3±0.2

**pH**

7.10-7.50

**Cultural Response**

M1273: Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours.

<b>Organism</b>	<b>Inoculum (CFU)</b>	<b>Growth</b>	<b>Pigment</b>
<i>Escherichia coli</i> ATCC 25922	>=10 <sup>3</sup>	inhibited	
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	good-luxuriant	blue green
<i>Stenotrophomonas maltophilia</i> ATCC 13637	>=10 <sup>3</sup>	inhibited	

**Storage and Shelf Life**

Store below 30°C in tightly closed container and the prepared medium at 2-8°C and use as fresh as possible. Use before expiry date on the label.

**Reference**

- 1.Brown M. R. W. and Scott F. J. H., 1970, J. Clin. Pathol., 23:172.
- 2.International Organization for Standardization (ISO), Draft ISO/DIS 8360-1:1988.

Revision : 1 / 2011

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.