



## Fraser Broth Base, Modified

M1764

### Intended use

It is recommended for the selective enrichment of *Listeria* species from foods.

### Composition\*\*

Ingredients	Gms / Litre
Peptone	5.000
Tryptone	5.000
Yeast extract	5.000
HM Peptone B #	5.000
Sodium chloride	20.000
Lithium chloride	3.000
Disodium phosphate	9.600
Monopotassium phosphate	1.350
Esculin	1.000
Nalidixic acid	0.010
Acriflavin	0.0125
Final pH ( at 25°C)	7.2±0.2

\*\*Formula adjusted, standardized to suit performance parameters

Key : # - Equivalent to Beef extract

### Directions

Suspend 54.97 grams in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add rehydrated contents of 2 vials of Fraser Supplement (FD141). Mix well and dispense as desired.

### Principle And Interpretation

*Listeria* species are widely distributed and are isolated from soil, decaying vegetable matter, sewage, water, animal feed, fresh and frozen poultry, meats, raw milk, cheese and asymptomatic human and animal carriers (1). Only *Listeria monocytogenes* from the genus *Listeria* ; causes infections in humans. *L.monocytogenes* primarily causes meningitis, encephalitis or septicemia in humans (2, 3). In pregnant women, *Listeria monocytogenes* often causes an influenza like bacteremic illness that, if untreated, may lead to amnionitis and infection of the fetus, resulting in abortion, still birth or premature birth. Contaminated foods are the primary vehicles of transmission (4).

Fraser Broth Base, Modified is based on the formulation by Fraser and Sperber (9). It is recommended for selective enrichment of *Listeria* species from foods.

This medium contains peptone, Tryptone, yeast extract and HM Peptone B which provide essential nutrients like carbon and nitrogenous compounds including vitamins, amino acids and trace ingredients. Phosphates buffer the medium while sodium chloride maintains osmotic equilibrium. Nalidixic acid and Acriflavin inhibits the growth of gram-negative and gram-positive organisms respectively (5,6,7) except *Listeria* species (5,6,7). *Listeria* species hydrolyze esculin to glucose and esculetin. The latter combines with ferric ions of ferric ammonium citrate (FD141), resulting in the formation of 6-7 dihydroxycoumarin, a black brown complex. Ferric ammonium citrate also enhances the growth of *L. monocytogenes* (8). The high salt tolerance (of sodium chloride) of *Listeria* is used as means to inhibit the growth of Enterococci. Lithium chloride is also used to inhibit Enterococci, which also possess the ability to hydrolyze esculin.

### Type of specimen

Food samples : meat and meat products; Dairy samples

### Specimen Collection and Handling:

For food and dairy samples, follow appropriate techniques for sample collection and processing as per guidelines (1,4,5).

After use, contaminated materials must be sterilized by autoclaving before discarding.

Please refer disclaimer Overleaf.

## Warning and Precautions :

Read the label before opening the container. Wear protective gloves/protective clothing/eye protection/ face protection. Follow good microbiological lab practices while handling specimens and culture. Standard precautions as per established guidelines should be followed while handling specimens. Safety guidelines may be referred in individual safety data sheets

## Limitations :

1. Due to variable nutritional requirements, some strains show poor growth on this medium.

## Performance and Evaluation

Performance of the medium is expected when used as per the direction on the label within the expiry period when stored at recommended temperature.

## Quality Control

### Appearance

Cream to yellow homogeneous free flowing powder

### Colour and Clarity of prepared medium

Fluorescent yellow coloured clear solution.

### Reaction

Reaction of 5.5% w/v aqueous solution at 25°C. pH : 7.2±0.2

### pH

7.00-7.40

### Cultural response

Cultural characteristics observed on addition of FD141 after an incubation at 35 - 37°C for 24-48 hours.

### Cultural Response

Organism	Inoculum (CFU)	Growth	Esculin Hydrolysis
<b>Cultural response</b>			
<i>Escherichia coli</i> ATCC 25922 (00013*)	≥10 <sup>3</sup>	inhibited	
<i>Enterococcus faecalis</i> ATCC 29212 (00087*)	50-100	none-poor	
<i>Listeria monocytogenes</i> ATCC 19111 (00020*)	50-100	good-luxuriant	positive, reaction, blackening of medium
<i>Listeria monocytogenes</i> ATCC 19112	50-100	good-luxuriant	positive reaction, blackening of medium
<i>Listeria monocytogenes</i> ATCC 19117	50-100	good-luxuriant	positive reaction, blackening of medium
<i>Listeria monocytogenes</i> ATCC 19118	50-100	good-luxuriant	positive reaction, blackening of medium
<i>Staphylococcus aureus</i> subsp. <i>aureus</i> ATCC 25923 (00034*)	50-100	none-poor	

Key : \* - Corresponding WDCM numbers

## Storage and Shelf Life

Store between 10-30°C in a tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label. On opening, product should be properly stored dry, after tightly capping the bottle in order to prevent lump formation due to the hygroscopic nature of the product. Improper storage of the product may lead to lump formation. Store in dry ventilated area protected from extremes of temperature and sources of ignition Seal the container tightly after use. Use before expiry date on the label.

Product performance is best if used within stated expiry period.

## Disposal

User must ensure safe disposal by autoclaving and/or incineration of used or unusable preparations of this product. Follow established laboratory procedures in disposing of infectious materials and material that comes into contact with sample must be decontaminated and disposed of in accordance with current laboratory techniques (6,7).

## Reference

1. Seeliger H. P. R., and Jones D., 1986, Bergeys Manual of Systematic Bacteriology, Vol. The Williams and Wilkins Co., Baltimore.
2. Nieman R. E., and Lorber B., 1980, Rev. Infect. Dis. 2 : 207-227.
3. Schuchat A. B., Swaminathan and C. V. Broome, Clin. Microbiol. Rev. 4: 169-183.
4. Murray P. R., Baron E. J., Jorgensen J. H., Tenover F. C., Tenover P. C., (Eds.), 8th Ed., 2003, Manual of Clinical Microbiology, ASM, Washington, D.C.
5. Lovette J., Francis D.W. and Hunt J.M., 1987, J. Food Prot., 50:188.
6. Lee W.K. and McClain D., 1986, Appl. Environ. Microbiol., 52:1215.
7. McClain D. and Lee W.H., 1988, J. Assoc. Off. Anal. Chem., 71:660.
8. Cowart R. E. and Foster B. G., 1985, J. Infect. Dis.; 151:172.
9. Fraser, J., and W. Sperber. 1988. Rapid detection of Listeria in food and environmental samples by esculin hydrolysis. Journal of Food Protection 51: 762-765.

Revision : 03 / 2018

### Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.