



## D.T.M. Agar Base (Dermatophyte Test Agar Base)

M188

Dermatophyte Test Medium (DTM) is used for selective isolation of dermatophytes.

### Composition\*\*

Ingredients	Gms / Litre
Papaic digest of soyabean meal	10.000
Glucose	10.000
Phenol red	0.200
Agar	20.000
Final pH ( at 25°C)	5.5±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Directions

Suspend 20.10 grams in 500 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 50°C. Aseptically add the rehydrated contents of one vial of Dermato Supplement (FD015). Mix well before pouring into sterile Petri plates.

### Principle And Interpretation

The Dermatophytes are a distinct group of fungi that infect the hair, skin and nails of humans and animals producing a variety of cutaneous infections known as ringworm (2). Dermatophytes like *Trichophyton*, *Microsporum* and *Epidermatophyton* are responsible for most of the cutaneous fungal infections (1). DTM Agar Base was developed by Taplin as a selective and differential medium for detection and identification of dermatophytes (2). On this medium identification of Dermatophytes are based on morphology and alkaline metabolites production. A combination of three antimicrobial agents (cycloheximide, chlortetracycline and gentamicin) inhibits bacteria and saprophytic yeasts and moulds. Dermatophytes are presumptively identified based on gross morphology and the production of alkaline metabolites, which raise the pH and cause the phenol red indicator to change the color of the medium from yellow to pink-red (2-4).

Papaic digest of soyabean meal provides nitrogenous and carbonaceous substances essential for growth. Glucose is the energy source. The pH indicator, phenol red, is used to detect amine production. Cycloheximide (5) (as FD) inhibits most of the saprophytic fungi. Gentamicin inhibits gram-negative bacteria including *Pseudomonas* species while chlortetracycline inhibits a wide range of gram-positive and gram-negative bacteria. The presence of growth on the medium provides presumptive identification of dermatophytes. D.T.M. Agar helps in isolation and early recognition of members of the *Microsporum*, *Trichophyton* by means of the distinct colour change from yellow to red. Rapidly growing species may effect a complete colour change within 3 days while slow growers will change colour in proportionately longer time. Non-Dermatophytes can be recognized by the absence of colour change. A few saprophytes, yeasts and bacteria change the medium from yellow to red, but can be easily distinguished by colonial morphology. Complete classification of Dermatophytes depends on microscopic observations along with biochemical and serological tests.

### Quality Control

#### Appearance

Light yellow to pink homogeneous free flowing powder

#### Gelling

Firm, comparable with 2.0% agar gel.

#### Colour and Clarity of prepared medium

Orange red coloured, slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 4% w/v aqueous solution at 25°C. pH : 5.5±0.2

**pH**

5.30-5.70

**Cultural Response**

M188: Cultural characteristics observed with added Dermato Supplement (FD015), after an incubation at 25-30°C for 6 days.

<b>Organism</b>	<b>Growth</b>	<b>Colour of Medium</b>
-----------------	---------------	-------------------------

**Cultural Response**

* <i>Aspergillus brasiliensis</i> ATCC 16404	none-poor	
---	-----------	--

<i>Candida albicans</i> ATCC 10231	good	
---------------------------------------	------	--

<i>Microsporum audouinii</i> ATCC 9079	good	pink-red
---	------	----------

<i>Pseudomonas aeruginosa</i> ATCC 27853	none-poor	
---	-----------	--

<i>Trichophyton mentagrophytes</i> ATCC 9533	good	pink-red
--	------	----------

Key : \*- Formerly known as *Aspergillus niger*

**Storage and Shelf Life**

Store below 30°C in tightly closed container and the prepared medium at 2-8°C. Use before expiry date on the label

**Reference**

1. Isenberg (Eds.), 1992, Clinical Microbiology Procedures Handbook, Vol . 1, American Society for Microbiology, Washington, D.C.
2. Taplin, Zaias, Rebell and Blank, 1969, Arch. Dermatol., 99:203-209.
3. Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Eds.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
4. Kwon-Chung and Bennett, 1992, Medical Mycology, Lea & Febiger, Philadelphia, Pa.
5. Rosenthal S., Stritzler R. and Villafane J., 1968, Arch. Dermatol., 97:685.

Revision : 1 / 2011

**Disclaimer :**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.