



FDA Agar (Extract Agar)

M236

FDA Agar (Extract Agar) is used for general cultivation of bacteria as well as routine testing of disinfectants and antiseptics.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	10.000
Beef extract	5.000
Sodium chloride	5.000
Agar	15.000
Final pH (at 25°C)	7.3±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 35 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

FDA Agar is used for general cultivation of bacteria as well as for routine testing of antiseptics and disinfectants. FDA Agar is also known as AATCC bacteriostasis agar where AATCC stands for American Association of Textile Chemists and Colourists (1). FDA agar is the formulation specified by Food and Drug Administration, U.S.A. and also by Association of Analytical Chemists (AOAC) (1, 2). It is used for detecting antibacterial activity of fabrics. FDA agar is a relatively simple formulation. Beef extract and peptic digest of animal tissue provide the nutrients required for microbial growth. Sodium chloride maintains osmotic equilibrium.

Quality Control

Appearance

Off white to yellow coloured homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Amber coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 3.5% w/v aqueous solution at 25°C. pH : 7.3±0.2

pH

7.10-7.50

Cultural Response

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Cultural Response

Organism	Inoculum (CFU)	Growth	Recovery
Cultural Response <i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	≥70%
<i>Staphylococcus aureus</i> ATCC 25923	50-100	good-luxuriant	≥70%
<i>Salmonella Typhi</i> ATCC 6539	50-100	good-luxuriant	≥70%

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. Technical Manual of AATCC, Vol. 61, 1985-86, AATCC, Research Triangle Park, N.C.
2. Williams (Ed.), 1984, Official Methods of Analysis of the AOAC, 14th ed. AOAC, Washington, D.C.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.