



E E Broth, Mossel

M287

E E Broth, Mossel is used for selective enrichment of *Enterobacteriaceae* in bacteriological examination of foods.

Composition**

| Ingredients | Gms / Litre |
|--------------------------------|-------------|
| Peptic digest of animal tissue | 10.000 |
| Dextrose | 5.000 |
| Disodium phosphate | 6.450 |
| Monopotassium phosphate | 2.000 |
| Ox bile, purified | 20.000 |
| Brilliant green | 0.0135 |
| Final pH (at 25°C) | 7.2±0.2 |

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 43.46 grams in 1000 ml distilled water. Dispense in tubes or flasks as desired. Stopper with cotton plugs or loose fitting caps. Heat in free flowing steam or boiling water for 30 minutes. Avoid overheating of the medium. DO NOT AUTOCLAVE.

Principle And Interpretation

The family *Enterobacteriaceae* consists of *Salmonella*, *Shigella* and other enteric pathogens. These organisms find entry into the foodsystem through faecally contaminated water. Majority of these organisms may be eliminated under the stringent food processing parameters. But some of these organisms may become sublethally injured during the changes in pH, exposure to steam or heat and other unfavourable conditions (1). Therefore the important aspect of food monitoring depends upon the identification and enumeration of these injured cells, after resuscitation. EE Broth, Mossel, formulated by Mossel et al (2) is recommended as an enrichment medium for *Enterobacteriaceae* in the biological examination of foods (2) and animal feed stuffs (3).

Peptic digest of animal tissue and dextrose provide the essential nutrients required for the growth of most of the members of *Enterobacteriaceae*. Brilliant green and ox bile, purified inhibit growth of gram-positive bacteria. Lactose-negative, anaerogenic lactose-positive or late lactose-fermenting *Enterobacteriaceae* are often missed by the standard coli-aerogenes test. To overcome this problem, lactose is replaced by dextrose in these media. Phosphates form the buffering system of the medium. The cells damaged while drying or low pH are resuscitated in well-aerated Tryptone Soya Broth (M011) for 2 hours at 25°C prior to enrichment in EE Broth. The resuscitation procedure is recommended for dried foods (5), animal feeds (6) and semi-preserved foods (7). EE Broth is an enrichment broth and should be used in conjunction with Violet Red Bile Glucose Agar (M581). Subcultures must be made onto lactose differential media such as MacConkey Agar (M081), Deoxycholate Citrate Agar (M065) or Brilliant Green Agar (M016) for the detection of lactose negative or delayed lactose fermenters. This is used to inoculate MPN tubes prepared using EE Broth. Inoculate a loopful from these tubes onto Violet Red Bile Glucose Agar (M581) after an initial incubation at 35-37°C for 24 hours. Typical pink colonies from M581 are subcultured for biochemical confirmation by oxidase and fermentation reactions (4). Decimal dilutions of the food homogenate are used if the expected counts are high or else initial suspension is used. EE Broth, Mossel (M287).

Quality Control

Appearance

Light yellow to greenish yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Emerald green coloured, clear solution without any precipitate

pH of 4.35% w/v aqueous solution at 25°C.

pH

7.00-7.40

Cultural Response

Cultural response was observed after an incubation at 35-37°C for 24-48 hours.

Cultural Response

| Organism | Inoculum (CFU) | Growth | Acid |
|--|----------------|-----------|----------------------------------|
| Cultural Response | | | |
| <i>Escherichia coli</i> ATCC 8739 | 50 -100 | luxuriant | positive reaction, yellow colour |
| <i>Pseudomonas aeruginosa</i> ATCC 9027 | 50 -100 | luxuriant | - |
| <i>Staphylococcus aureus</i> ATCC 6538 | $\geq 10^3$ | inhibited | |
| <i>Escherichia coli</i> ATCC 25922 | 50 -100 | luxuriant | positive reaction, yellow colour |
| <i>Escherichia coli</i> NCTC 9002 | 50 -100 | luxuriant | positive reaction, yellow colour |
| <i>Pseudomonas aeruginosa</i> ATCC 27853 | 50 -100 | luxuriant | - |
| <i>Enterobacter aerogenes</i> ATCC 13048 | 50 -100 | luxuriant | positive reaction, yellow colour |
| <i>Proteus mirabilis</i> ATCC 25933 | 50 -100 | luxuriant | positive reaction, yellow colour |
| <i>Salmonella Enteritidis</i> ATCC 13076 | 50 -100 | luxuriant | variable reaction |
| <i>Shigella boydii</i> ATCC 12030 | 50 -100 | luxuriant | negative reaction |
| <i>Staphylococcus aureus</i> ATCC 25923 | $\geq 10^3$ | inhibited | |

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1. Mossel D. A. A., and Harrewijn G. A., 1972, Alimenta II, 29-30
2. Mossel D. A. A., Vissar M. and Cornellsen A. M. R., 1963, J. Appl. Bacteriol., 26(3):444.
3. Van Schothurst M. et al, 1966, Vet Med., 13(3):273.
4. International Organization for Standardization (ISO), 1993, Draft ISO/DIS 7402.
5. Mossel D. A. A. and Ratto M. A., 1970, Appl. Microbiol., 20:273.
6. Mossel D. A. A., Shennan J. L. and Clare V., 1973, J. Sci. Fd. Agric., 24: 499.
7. Mossel D. A. A. and Ratto M. A., 1973, J. Food Technol., 8: 97.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.