



Sulphate API Broth w/o Sodium Lactate

M310

Sulphate API Broth is used for detection and estimation of sulphate reducing bacteria.

Composition**

Ingredients	Gms / Litre
Yeast extract	1.000
Magnesium sulphate	0.200
Dipotassium phosphate	0.010
Ferrous ammonium sulphate	0.100
Sodium chloride	10.000
Ascorbic acid	0.10
Final pH (at 25°C)	7.5±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 11.41 grams in 1000 ml distilled water. Add 4 ml of sodium lactate. Heat if necessary to dissolve the medium completely. Dispense, preferably in screw-capped tubes in 9 ml amounts. Sterilize by autoclaving at 15 lbs pressure (121°C) for 10 minutes. Close the caps immediately while the medium is still hot.

Principle And Interpretation

Sulphate API Broth is prepared according to the formulation described in the American Petroleum Institute Recommended Practice (1) for detection of sulphate reducing bacteria. Sulphate-reducing bacteria cause corrosion of oil well systems resulting in perforations in the pipes. Sulphate-reducing bacteria convert sulphate to sulphide which with the ferrous ion gives black colour. The insoluble sulphide results in plugging.

Yeast extract in the medium provides nitrogen and other nutrients necessary to support bacterial growth. Ascorbic acid is the carbohydrate source. Dipotassium Phosphate buffers the medium. Sodium chloride, magnesium sulphate and ferrous ammonium sulphate provide essential ions. *Desulfovibrio* oxidizes reduced substrates i.e. sodium lactate, further with stepwise reduction of sulfate to sulfide. The detection and estimation of these bacteria is done on the basis of their ability to grow and produce sulphide in this medium. For the estimation, appropriate dilutions of water samples are inoculated.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured clear solution in tubes

Reaction

Reaction of 1.1% w/v aqueous solution (containing 0.4% v/v Sodium lactate) at 25°C. pH : 7.5±0.2

pH

7.30-7.70

Cultural Response

Cultural characteristics observed after an incubation at 30°C for upto 1 week, under anaerobic condition.

Cultural Response

Organism	Inoculum (CFU)	Growth
Cultural Response <i>Desulfovibrio desulfuricans</i> <i>ATCC 13541</i>	50-100	good-luxuriant

Storage and Shelf Life

Store dehydrated medium and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Reference

1.American Petroleum Institute Recommended Practice 28, 1959, First ed.

Revision : 02/ 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory,diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.