



Actinomyces Agar

M341

Actinomyces Agar is recommended for the cultivation and maintenance of the anaerobic *Actinomyces* species.

Composition**

Ingredients	Gms / Litre
Beef heart infusion, solids	10.000
Tryptose	10.000
Casein enzymic hydrolysate	4.000
Yeast extract	5.000
Dextrose	5.000
L-Cysteine hydrochloride	1.000
Starch, soluble	1.000
Sodium chloride	5.000
Monopotassium phosphate	15.000
Ammonium sulphate	1.000
Magnesium sulphate	0.200
Calcium chloride	0.020
Agar	20.000
Final pH (at 25°C)	6.9±0.2

**Formula adjusted, standardized to suit performance parameters

Directions

Suspend 77.22 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Distribute into tubes or flasks. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Principle And Interpretation

Actinomycetes are gram-positive bacteria, which show marked chemical and morphological diversity but form a distinct evolutionary line of organisms that range from coccoid and pleomorphic forms to branched filaments (1). *Actinomycetes* form an integral part of soil, water and vegetation. *Actinomycete* development leads to the formation of volatile metabolites (2). Traces of these volatile metabolites are sufficient to impart disagreeable odour to water or a muddy flavour to fish (3). *Actinomycetes* also cause disruptions in wastewater treatment by forming massive growths, which are capable of producing thick foam in the activated sludge process (4, 5) Actinomyces Agar/Broth is further modified and is recommended for the cultivation and maintenance of anaerobic *Actinomycete* species (6).

.Beef heart infusion, tryptose, casein enzymic hydrolysate, yeast extract, starch and dextrose act as sources of carbon, nitrogen, sulphur, vitamins and other growth factors. The metallic salts provide essential electrolytes and minerals.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 2.0% agar gel.

Colour and Clarity of prepared medium

Yellow to light amber coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 7.72% w/v aqueous solution at 25°C. pH : 6.9±0.2

pH

6.70-7.10

Cultural Response

M341: Cultural characteristics observed after an incubation at 25-30°C for 40-72 hours (*- incubated anaerobically)

Organism	Growth
Cultural Response	
* <i>Actinomyces israelii</i> ATCC 10049	luxuriant
<i>Streptomyces achromogenes</i> ATCC 12767	good
<i>Streptomyces albus</i> subsp <i>albus</i> ATCC 3004	good
<i>Streptomyces lavendulae</i> ATCC 8664	good
* <i>Actinomyces bovis</i> ATCC 13683	good

Storage and Shelf Life

Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on label.

Reference

1. Collee J. G., Fraser A. G., Marmion B. P., Simmons A., (Eds.), Mackie and McCartney, Practical Medical Microbiology, 1996, 14th Edition, Churchill Livingstone.
2. Adams B. A., 1929, Water and Water Eng., 31:327.
3. Eaton A. D., Clesceri L. S. and Greenberg A. W., (Eds.), 2005, Standard Methods for the Examination of Water and Wastewater, 21st Ed., APHA, Washington, D.C.
4. Lechevalier H. A., 1975, Environ. Protection Technol. Ser., EPA-600/ 2-75-031, U. S. Environmental Protection Agency, Cincinnati, Ohio.
5. Lechevalier M. P., and Lechevalier H. A., 1974, Int. J. Syst.Bacteriol., 24:278.
6. Ajello L., Georg L. K., Kaplan W. and Kaufman L., 1963, CDC Lab Manual Med. Mycology, PHS Publication No. 994, CDC, Washington D.C.

Revision : 2 / 2015

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related HiMedia™ publications. The information contained in this publication is based on our research and development work and is to the best of our knowledge true and accurate. HiMedia™ Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.